



UNIVERSIDAD DE
MONTEVIDEO

FACULTAD DE CIENCIAS EMPRESARIALES Y ECONOMIA

**Serie de documentos de trabajo del Departamento de Economía /
Department of Economics Working Papers Series**

CONTINUITY AND COMPLETENESS UNDER RISK*

Juan Dubra
Departamento de Economía
Facultad de Ciencias Empresariales y Economía
Universidad de Montevideo
jdubra@um.edu.uy

Working paper UM_CEE_2010-04
<http://www.um.edu.uy/cee/investigaciones/>

The working papers of the Department of Economics, Universidad de Montevideo are circulated for discussion and comment purposes. They have not been peer reviewed nor been subject to the review by the University's staff.

© 2010 by Juan Dubra. All rights reserved. Short sections of text, not to exceed two paragraphs, may be quoted without explicit permission provided that full credit, including © notice, is given to the source.

Continuity and Completeness under Risk*

Juan Dubra

Universidad de Montevideo

December 2, 2010

Let X be a finite set, and for $m = |X|$, let $\mathcal{P} = \{p \in \mathbf{R}_+^m : \sum_i p_i = 1\}$ be the set of lotteries over X . Let \succeq be a transitive and reflexive binary relation on \mathcal{P} . As usual define $s \succ t$ if $s \succeq t$ and not $t \succeq s$, and $s \sim t$ if $s \succeq t$ and $t \succeq s$. We say that \succeq is non trivial if there exist s and t in \mathcal{P} such that $s \succ t$. The relation \succeq satisfies:

Independence, if for all $p, q, r \in \mathcal{P}$ and $\lambda \in (0, 1)$, $p \succeq q$ if and only if $\lambda p + (1 - \lambda)r \succeq \lambda q + (1 - \lambda)r$;

Herstein Milnor, if for all $p, q, r \in \mathcal{P}$ the set $\{\alpha \in [0, 1] : \alpha p + (1 - \alpha)q \succeq r\}$ is closed;¹

Archimedean, if for all $p, q, r \in \mathcal{P}$, $p \succ q$ implies $\lambda p + (1 - \lambda)r \succ q$ for some $\lambda \in (0, 1)$;

Completeness, if for all p and q , either $p \succeq q$ or $q \succeq p$.

In this note I prove the following Theorem.

Theorem 1 *Suppose \succeq is a transitive, reflexive, non-trivial binary relation on \mathcal{P} , that satisfies Independence. If \succeq satisfies any two of the following axioms, it satisfies the third: Herstein-Milnor, Archimedean and Completeness.*

Schmeidler (1971) proved an analogous theorem for the case in which \succeq is a preference relation on a set, not necessarily involving lotteries. He proved that if \succeq on a connected topological set Z is such that for some x and y , $x \succ y$, then closed weak upper and lower contour sets and open strict upper and lower contour sets imply completeness.

That Completeness and Independence imply that HM Continuity and Archimedean are equivalent is trivial and was first claimed by Aumann (1962, p. 453). Karni (2007) proved that under a property weaker than Independence (Local Mixture Dominance), Completeness and Archimedean imply HM Continuity. Hence, we only need to prove that under the assumptions of the Theorem, HM Continuity and Archimedean imply Completeness; to do so I will prove the following Lemma, which together with Schmeidler's theorem will establish the desired result.

Lemma 1 *Suppose X is finite, that \succeq is a transitive, reflexive binary relation on the space \mathcal{P} of lotteries over X , and that \succeq satisfies Independence.*

*I thank Edi Karni, Efe Ok, and a referee in this journal for comments.

¹The proof below shows that under Independence, this version of HM implies the stronger version which also requires that $\{\alpha : r \succeq \alpha p + (1 - \alpha)q\}$ is closed. A similar argument applies to the definition of the Archimedean axiom.

- a) If \succeq satisfies HM Continuity then for all p , $\{q : q \succeq p\}$ and $\{q : p \succeq q\}$ are closed.
- b) If \succeq satisfies the Archimedean Axiom, then for all p , $\{q : q \succ p\}$ and $\{q : p \succ q\}$ are open in the relative topology in \mathcal{P} .

A version of part (a) of the Lemma was established in Proposition 1 in Dubra et al. (2004), but with slightly different axioms: a weaker Independence, and a stronger continuity: Double Mixture Continuity: for any p, q, r, s in \mathcal{P} the following set is closed

$$T = \{\lambda \in [0, 1] : \lambda p + (1 - \lambda)r \succeq \lambda q + (1 - \lambda)s\}.$$

Part (a) of the Lemma is relevant, despite Proposition 1 in Dubra et al., because Double Mixture Continuity is not a standard axiom, and the Independence axiom in this paper is standard. Also, the proof is similar, but simpler. To the best of my knowledge, part (b) is new. Both (a) and (b) could be proved in a more cumbersome manner by appealing to the well known equivalence between algebraic closedness (HM Continuity) and topological closedness (and similarly for openness).

Proof. Proof of (a). In order to show that for all v the set $S = \{r : r \succeq v\}$ is closed take any q in its boundary. If S is a singleton, there is nothing to prove, and if it is not, by the Independence axiom it is a convex set and therefore has a nonempty relative interior. Pick any p in the relative interior of S .

Let B be the open unit ball in the linear space generated by $S - v$, endowed with the relative topology. Fix any $\lambda \in (0, 1)$ and any $\varepsilon > 0$. For any $b \in B$, pick $\delta > 0$ small enough that $\varepsilon b + \delta B \subset \varepsilon B$. Since q is in the boundary of S , there exists $w \in S$ such that $\|w - q\| < \delta$, which implies $\varepsilon b + (1 - \lambda)(q - w) \in \varepsilon B$ and therefore

$$\begin{aligned} \lambda p + (1 - \lambda)q + \varepsilon b &= \lambda p + (1 - \lambda)w + (1 - \lambda)(q - w) + \varepsilon b \\ &\in \lambda p + (1 - \lambda)S + \varepsilon B. \end{aligned} \tag{1}$$

For fixed λ , since p is in the relative interior of S , there exists $\varepsilon > 0$ small enough such that $p + \frac{\varepsilon}{\lambda}B \subseteq S$. Since equation (1) was true for all ε , we obtain

$$\lambda p + (1 - \lambda)q + \varepsilon B \subseteq \lambda \left(p + \frac{\varepsilon}{\lambda}B \right) + (1 - \lambda)S \subseteq \lambda S + (1 - \lambda)S = S.$$

Then, since $0 \in B$, we get that for all $\lambda \in (0, 1)$, $\lambda p + (1 - \lambda)q \in S$, and by HM, $q \in S$, as was to be shown.

Consider now lower contour sets: for some fixed v , let $T = \{r : v \succeq r\}$. By Independence, for $u = (\frac{1}{m}, \dots, \frac{1}{m})$, $v \succeq r$ if and only if $v - r \in \{\lambda(p - u) : p \succeq u, \lambda > 0\}$. Hence

$$T = \{r : v \succeq r\} = \{v - m(p - u) : p \succeq u\} \cap \mathcal{P},$$

and closedness follows by closedness of $\{p : p \succeq u\}$.

Proof of (b). We will show that for all p , $\{q : q \succ p\}$ and $\{q : p \succ q\}$ are relatively open in \mathcal{P} by showing that $D = \{\lambda(r - u) : \lambda > 0 \text{ and } r \succ u\}$ is relatively open in the linear space generated by $\mathcal{P} - u$, which we denote $A \equiv \text{aff}(\mathcal{P} - u)$. Openness of D implies openness of $\{q : q \succ p\} = (p + D) \cap \mathcal{P}$ (where the equality follows by Independence). Openness of D also implies openness of $\{q : p \succ q\} = (p - D) \cap \mathcal{P}$.

It is easy to see that $A = \{\mu \in \mathbf{R}^m : \sum_1^m \mu_i = 0\}$. Also, given any $\mu \in A$, one can pick λ large enough so as to make $\max_{i \leq m} |\mu_i| < \frac{\lambda}{m}$. Define then $p_i = \frac{\mu_i}{\lambda} + \frac{1}{m}$; it is then straightforward to check that $p \in \mathcal{P}$ and that $\mu = \lambda(p - u)$, showing $A = \{\lambda(p - u) : \lambda > 0, p \in \mathcal{P}\}$.

To show that D is relatively open, pick any $\sigma \in D$ and let $\sigma = \lambda'(p' - u) \in D$. For small enough α , $p \equiv \alpha p' + (1 - \alpha)u$ is in the relative interior of \mathcal{P} and by Independence, $p' \succ u$ ensures that $p \succ u$. Let $\lambda = \frac{\lambda'}{\alpha}$ and note that

$$\sigma = \lambda'(p' - u) = \frac{\lambda'}{\alpha} \alpha(p' - u) = \lambda \alpha(p' - u) = \lambda(p - u).$$

For $i = 1, \dots, m-1$, let $c_i = e_i - e_m$ be a basis for A , and for $i = m, \dots, 2m-2$, let $c_i = -c_{i-m+1}$. Since p is in the interior of \mathcal{P} , for small enough α_i , $p + \alpha_i c_i \in \mathcal{P}$, and since $p \succ u$, the Archimedean axiom ensures that there is some β_i such that $p + \beta_i \alpha_i c_i = (1 - \beta_i)p + \beta_i(p + \alpha_i c_i) \succ u$. Using Independence again, we obtain that for $\gamma_i \equiv \beta_i \alpha_i$ and $\gamma \equiv \min_i \gamma_i > 0$,

$$p + \gamma c_i = \frac{\gamma}{\gamma_i} (p + \gamma_i c_i) + \left(1 - \frac{\gamma}{\gamma_i}\right) p \succ u \quad (2)$$

for all i . Let H denote the convex hull of $\{\gamma c_i\}_{i=1}^{2m-2}$, and note that since for each i , H contains both c_i and $-c_i$,

$$\text{if } \pi \in H \text{ and } \delta < 1 \text{ then } \delta \pi \in H. \quad (3)$$

Moreover, by (2) and Independence, for any s ,

$$s \in p + H \Rightarrow s \succ u. \quad (4)$$

For any $x, y \in \mathbf{R}^m$ let $d(x, y) = \max_i |x_i - y_i|$, and let $B = \{\mu \in A : d(\mu, 0) < \frac{\gamma}{m-1}\}$ be a relatively open set in A . Notice that for any $\mu \in B$, since $\mu = \sum_{i=1}^{m-1} \mu_i c_i$ and $|\mu_i| < \frac{\gamma}{m-1}$, we obtain that for $\delta \equiv \sum_{i=1}^{m-1} \frac{|\mu_i|}{\gamma} < 1$, $\sum_{i=1}^{m-1} \frac{|\mu_i|}{\gamma \delta} = 1$ and therefore

$$\mu = \sum_{i \leq m-1: \mu_i > 0} \mu_i c_i + \sum_{i \leq m-1: \mu_i < 0} (-\mu_i) (-c_i) = \delta \left[\sum_{i: \mu_i > 0} \frac{\mu_i}{\gamma \delta} \gamma c_i + \sum_{i: \mu_i < 0} \frac{-\mu_i}{\gamma \delta} (-\gamma c_i) \right]$$

showing that $\mu = \delta \pi$ for π a convex combination of $\{\gamma c_i\}_{i=1}^{2m-2}$, and $\delta < 1$. By equation (3), μ is in H .

For any $\sigma' \in \sigma + \lambda B = \lambda(p - u) + \lambda B$, we have that for some $\mu \in B \subseteq H$, $u + \frac{\sigma'}{\lambda} = p + \mu \in p + B \subseteq p + H$. By equation (4) we obtain $u + \frac{\sigma'}{\lambda} \succ u$ which implies $\sigma' \in D$, which shows that D is relatively open since for any $\sigma \in D$ we have found an open set $\sigma + \lambda B$ containing σ such that $\sigma + \lambda B \subset D$. ■

References

- Aumann, R. (1962), "Utility theory without the completeness axiom," *Econometrica* 30(3).
 Dubra, J., F. Maccheroni and E. Ok (2004) "Expected Utility Theory without the Completeness Axiom," *Journal of Economic Theory* 115(1).
 Karni, E. (2007), "Archimedean and Continuity," *Mathematical Social Sciences* 53(3).
 Schmeidler, D. (1971), "A Condition for the Completeness of Partial Preference Relations," *Econometrica* 39(2).